

On the Irreducibility of Commuting Varieties of Nilpotent Matrices

Roberta Basili

*Dipartimento di Matematica e Informatica, Università di Perugia,
Via Vanvitelli 1, 06123 Perugia, Italy, e-mail: basili@dipmat.unipg.it*

Abstract

Given an $n \times n$ nilpotent matrix over an algebraically closed field K , we prove some properties of the set of all the $n \times n$ nilpotent matrices over K which commute with it. Then we give a proof of the irreducibility of the variety of all the pairs (A, B) of $n \times n$ nilpotent matrices over K such that $[A, B] = 0$ if either $\text{char } K = 0$ or $\text{char } K \geq \frac{n}{2}$. We get as a consequence a proof of the irreducibility of the local Hilbert scheme of n points of a smooth algebraic surface over K if either $\text{char } K = 0$ or $\text{char } K \geq \frac{n}{2}$.

1 Introduction

Let K be an algebraically closed field. T.S. Motzkin and O. Taussky in [9] and M. Gerstenhaber in [6] proved that for any $n \in \mathbb{N}$ the variety of all the pairs (A, B) of $n \times n$ matrices over K such that $[A, B] = 0$ is irreducible. R.W. Richardson in [12] extended this result by showing that if $\text{char } K = 0$ the variety of all the pairs of commuting elements of a reductive Lie algebra over K is irreducible. D.I. Panyushev in [11] studied this kind of varieties in a more general context and, besides other results, showed that if $\text{char } K = 0$ the variety of all the pairs of commuting elements of a symmetric space of maximal rank is normal (see also [3]).

Let $\mathcal{H}(n, K)$ be the variety of all the pairs (A, B) of $n \times n$ nilpotent matrices over K such that $[A, B] = 0$. Recently V. Baranovsky proved in [1] that $\mathcal{H}(n, K)$ is irreducible if either $\text{char } K = 0$ or $\text{char } K > n$. The proof uses the irreducibility of the local punctual Hilbert schemes of smooth algebraic surfaces. In fact, let $\mathcal{X}$ be an algebraic surface over K and let $\text{Hilb}^n \mathcal{X}$ be the Hilbert scheme of n points in $\mathcal{X}$. Let P be a nonsingular point of $\mathcal{X}$

and let $\text{Hilb}^n(\mathcal{O}_P)$ be the fiber in nP of the Hilbert-Chow morphism from $\text{Hilb}^n \mathcal{X}$ to $\text{Sym}^n(\mathcal{X})$. It parametrizes the ideals of colength n of $\mathcal{O}_P$. Let $\widehat{\mathcal{H}}(n, K)$ be the subvariety of $\mathcal{H}(n, K) \times K^n$ of all the triples (A, B, v) such that $\dim \langle A^i B^j v : i, j = 0, \dots, n-1 \rangle = n$. H. Nakajima in [10] showed that there exists a morphism from $\widehat{\mathcal{H}}(n, K)$ onto $\text{Hilb}^n(\mathcal{O}_P)$ whose fibers are the orbits of $\widehat{\mathcal{H}}(n, K)$ with respect to the action of $\text{GL}(n, K)$. If x, y are local coordinates in P and we regard $\mathcal{O}_P$ as a subset of $K[[x, y]]$, it associates to (A, B, v) the ideal of all $g \in \mathcal{O}_P$ such that $g(A, B)v = 0$. J. Briançon in [4] proved that $\text{Hilb}^n(\mathcal{O}_P)$ is irreducible if $\text{char } K = 0$ (see also [7]). In [8] this was extended by A.A. Iarrobino to the case $\text{char } K > n$.

In this paper we give a more elementary proof of the irreducibility of $\mathcal{H}(n, K)$ if either $\text{char } K = 0$ or $\text{char } K \geq \frac{n}{2}$. It consists of showing that the subset of $\mathcal{H}(n, K)$ of all (A, B) such that B is regular, that is $\text{rank } B = n-1$, is dense. For this purpose we fix an $n \times n$ nilpotent matrix B over K and study the Jordan canonical form of the $n \times n$ nilpotent matrices over K which commute with B . If A commutes with B we determine some properties of the Jordan canonical form of $B + f(A)$ for some suitable $f \in K[x]$ such that $x \mid f$. We use the form of the centralizer of B when B is in Jordan canonical form and the irreducibility of the variety of its nilpotent elements. The main step of the proof of the irreducibility of $\mathcal{H}(n, K)$ is the following result: if either $\text{char } K = 0$ or $\text{char } K \geq \frac{n}{2}$ and $m \in \mathbb{N}$ is such that $\frac{n}{2} + 2 < m \leq n$ the subvariety of $\mathcal{H}(n, K)$ of all the pairs (A, B) such that $\text{rank } A \geq n-2$ and $\frac{n}{2} + 1 < \text{ind } A \leq m$ (where $\text{ind } A$ is the index of nilpotency of A) is irreducible. In the proof of this result we use the same map introduced by Nakajima in [10] from an open subset of $\widehat{\mathcal{H}}(n, K)$ to the grassmannian of all the subspaces of $K[x, y]/(x, y)^n$ of codimension n . This result is not true for some $n, p \in \mathbb{N}$ such that $p < \frac{n}{2}$ and $\text{char } K = p$. As a consequence we get a proof of the irreducibility of $\text{Hilb}^n(\mathcal{O}_P)$ for algebraically closed fields K such that either $\text{char } K = 0$ or $\text{char } K \geq \frac{n}{2}$.

2 Preliminaries

If R is a ring and $n', n'' \in \mathbb{N}$ let $\mathcal{M}(n' \times n'', R)$, $\mathcal{M}(n', R)$ and $\mathcal{N}(n', R)$ be the varieties of all the $n' \times n''$ matrices, of all the $n' \times n'$ matrices and of all the $n' \times n'$ nilpotent matrices respectively over R . Let J_n be the nilpotent Jordan block of order n over K . In this section we will not use the hypothesis that K is algebraically closed.

Proposition 2.1 $\mathcal{N}(n, K)$ is irreducible of dimension $n^2 - n$.

Proof. If $\mathcal{N}_1(n, K)$ is the subspace of $\mathcal{N}(n, K)$ of the matrices whose entries of indices (i, j) such that $j - i \neq 1$ are 0, the morphism from $\text{GL}(n, K) \times \mathcal{N}_1(n, K)$ to $\mathcal{N}(n, K)$ defined by $(G, A) \mapsto G^{-1}AG$ is surjective. Since the centralizer of J_n has dimension n , the dimension of the orbit of J_n is $n^2 - n$, which shows the claim. $\square$

Let $B \in \mathcal{N}(n, K)$ be fixed, let $\mathcal{N}_B = \{A \in \mathcal{N}(n, K) : [A, B] = 0\}$ and let $u_1 \geq \dots \geq u_t$ be the orders of the Jordan blocks of B . We choose a basis $\Delta_B = \{v_h^{u_h-1}, \dots, v_h^0 : h = 1, \dots, t\}$ of K^n with respect to which B is in Jordan canonical form. If F is an extension field of K and $X \in \mathcal{M}(n, F)$ we regard the matrix which represents X with respect to Δ_B as a block matrix (X_{hk}) , $h, k = 1, \dots, t$, where $X_{hk} \in \mathcal{M}(u_h \times u_k, F)$. If $m \in \mathbb{N}$, $l, l' = 1, \dots, m$, and $(X_{ll'})$ is a block matrix, let $X_{(l)}$ and $X^{(l)}$ be respectively the l -th row and the l -th column of blocks of $(X_{ll'})$ for $l = 1, \dots, m$.

Lemma 2.2 (H.W. Turnbull and A.C. Aitken, 1931). *If $A \in \mathcal{M}(n, K)$ we have $[A, B] = 0$ if and only if for $1 \leq k \leq h \leq t$ the matrices A_{hk} and A_{kh} have the following form:*

$$A_{hk} = \begin{pmatrix} 0 & \dots & 0 & a_{hk}^1 & a_{hk}^2 & \dots & a_{hk}^{u_h} \\ \vdots & & & 0 & a_{hk}^1 & \ddots & \vdots \\ \vdots & & & & \ddots & \ddots & a_{hk}^2 \\ 0 & \dots & \dots & \dots & \dots & 0 & a_{hk}^1 \end{pmatrix},$$

$$A_{kh} = \begin{pmatrix} a_{kh}^1 & a_{kh}^2 & \dots & a_{kh}^{u_h} \\ 0 & a_{kh}^1 & \ddots & \vdots \\ \vdots & \ddots & \ddots & a_{kh}^2 \\ \vdots & & 0 & a_{kh}^1 \\ \vdots & & & 0 \\ \vdots & & & \vdots \\ 0 & \dots & \dots & 0 \end{pmatrix}$$

where for $u_h = u_k$ we omit the first $u_k - u_h$ columns and the last $u_k - u_h$ rows respectively.

Proof. See [14], [5], [2]. $\square$

Let $q_0 = 0$ and let $q_\alpha \in \{1, \dots, t\}$, $\alpha = 1, \dots, \hat{t}$, be such that $u_h = u_{h+1}$ if $q_{\alpha-1} + 1 \leq h < q_\alpha$, $u_{q_\alpha} \neq u_{q_\alpha+1}$, $q_{\hat{t}} = t$. If (A_{hk}) , $h, k = 1, \dots, t$, has the form of Lemma 2.2, for $\alpha, \beta = 1, \dots, \hat{t}$ let

$$\overline{A}_{\alpha\beta} = (a_{hk}^1), \quad q_{\alpha-1} + 1 \leq h \leq q_\alpha, \quad q_{\beta-1} + 1 \leq k \leq q_\beta.$$

Lemma 2.3 *If $A \in \mathcal{M}(n, K)$ is such that $[A, B] = 0$ then $A \in \mathcal{N}_B$ if and only if $\overline{A}_{\alpha\alpha}$ is nilpotent for $\alpha = 1, \dots, \hat{t}$, hence $\mathcal{N}_B$ is irreducible. Moreover if $A \in \mathcal{N}_B$ it is possible to choose Δ_B such that $\overline{A}_{\alpha\alpha}$ is upper triangular for $\alpha = 1, \dots, \hat{t}$.*

Proof. For $l = 0, \dots, u_1 - 1$ let $U^l = \langle v_h^l : h = 1, \dots, t, u_h - 1 \geq l \rangle$; then $K^n = \bigoplus_{l=0}^{u_1-1} U^l$ and $A(U^l) \subseteq \bigoplus_{i=l}^{u_1-1} U^i$. For $v \in K^n$ let $v = \sum_{l=0}^{u_1-1} v^{(l)}$ where $v^{(l)} \in U^l$ and let $L_{A,l} : U^l \rightarrow U^l$ be defined by $L_{A,l}(v) = A(v)^{(l)}$. Then A is nilpotent if and only if $L_{A,l}$ is nilpotent for $l = 0, \dots, u_1 - 1$. For $l = 0, \dots, u_1 - 1$ let $\gamma_l \in \{1, \dots, \hat{t}\}$ be such that $l \leq u_{q_{\gamma_l}} - 1$, $l > u_{q_{\gamma_l+1}} - 1$ if $\gamma_l \neq \hat{t}$. Then the matrix of $L_{A,l}$ with respect to the basis $\{v_h^l : h = 1, \dots, t, u_h - 1 \geq l\}$ is the lower triangular block matrix $(\overline{A}_{\alpha\beta})$, $\alpha, \beta = 1, \dots, \gamma_l$, which is nilpotent if and only if $\overline{A}_{\alpha\alpha}$ is nilpotent for $\alpha = 1, \dots, \gamma_l$. This proves the first claim and, by Proposition 2.1, the second one.

For $v \in K^n$ let $v = \sum_{\alpha=1}^{\hat{t}} v_{(\alpha)}$ where $v_{(\alpha)} \in \langle v_h^l : h = q_{\alpha-1} + 1, \dots, q_\alpha, l = 0, \dots, u_{q_\alpha} - 1 \rangle$. For $\alpha = 1, \dots, \hat{t}$ and $l = 0, \dots, u_{q_\alpha} - 1$ let $U_\alpha^l = \langle v_h^l : h = q_{\alpha-1} + 1, \dots, q_\alpha \rangle$ and let $L_{A,\alpha,l} : U_\alpha^l \rightarrow U_\alpha^l$ be defined by $L_{A,\alpha,l}(v) = L_{A,l}(v)_{(\alpha)}$. Then the matrix of $L_{A,\alpha,l}$ with respect to the basis $\{v_h^l : h = q_{\alpha-1} + 1, \dots, q_\alpha\}$ is $\overline{A}_{\alpha\alpha}$. Hence for $\alpha = 1, \dots, \hat{t}$ there exists $\{c(A, \alpha)_{hk} \in K : h, k = q_{\alpha-1} + 1, \dots, q_\alpha\}$ such that if we set

$$w_h^l = \sum_{k=q_{\alpha-1}+1}^{q_\alpha} c(A, \alpha)_{hk} v_k^l$$

for $h = q_{\alpha-1} + 1, \dots, q_\alpha$ we have that the matrix of $L_{A,\alpha,l}$ with respect to the basis $\{w_h^l : h = q_{\alpha-1} + 1, \dots, q_\alpha\}$ is upper triangular for $l = 0, \dots, u_{q_\alpha} - 1$. Then the basis $\{w_h^{u_h-1}, \dots, w_h^0 : h = 1, \dots, t\}$ has the required property. $\square$

We denote by $\mathcal{N}_{B,\Delta_B}$ the subspace of $\mathcal{N}_B$ of all A such that $\overline{A}_{\alpha\alpha}$ is upper triangular for $\alpha = 1, \dots, \hat{t}$.

Let $n_i \in \{1, \dots, t\}$, $i = 1, \dots, r_B$, be such that $n_1 = 1$, $u_{n_i} - u_{n_{i+1}} \geq 2$, $u_{n_i} - u_{n_{i+1}-1} \leq 1$ for $i = 1, \dots, r_B - 1$, $u_{n_{r_B}} - u_t \leq 1$.

Proposition 2.4 *There exists a non-empty open subset of $\mathcal{N}_B$ such that if A belongs to it we have $\text{rank } A = n - r_B$.*

Proof. By Lemma 2.3 it is sufficient to prove that there exists a non-empty open subset $\mathcal{Z}$ of $\mathcal{N}_{B,\Delta_B}$ such that $\text{rank } A = n - r_B$ for $A \in \mathcal{Z}$.

If $r_B = 1$ there exists $A \in \mathcal{N}_B$ regular such that B is a power of A and if $A \in \mathcal{N}_{B,\Delta_B}$ is regular then for $m = 1, \dots, u_1 - 1$ we have

$$v_1^{u_1-j}, \dots, v_t^{u_t-j} \in \langle Av_1^{u_1-j-1}, Av_2^{u_2-j}, \dots, Av_t^{u_t-j} : j = 1, \dots, m \rangle$$

for $j = 1, \dots, m$.

Let $r_B > 1$. For $i = 1, \dots, r_B$ and $A \in \mathcal{N}_{B,\Delta_B}$ we define $V_{A,i}$ to be the following subset of K^n :

$$\langle Av_{n_{i'}}^{u_{n_{i'}}-j-1}, Av_{n_{i'}+1}^{u_{n_{i'}+1}-j}, \dots, Av_{n_{i'}+1-1}^{u_{n_{i'}+1-1}-j} : j = 1, \dots, i'-i+1, i' = 1, \dots, i \rangle.$$

Then we have

$$V_{A,i} \subseteq \langle v_{n_{i'}}^{u_{n_{i'}}-j}, v_{n_{i'}+1}^{u_{n_{i'}+1}-j}, \dots, v_{n_{i'}+1-1}^{u_{n_{i'}+1-1}-j} : j = 1, \dots, i'-i+1, i' = 1, \dots, i \rangle$$

for $i = 1, \dots, r_B$. Moreover the open subset $\mathcal{Z}$ of $\mathcal{N}_{B,\Delta_B}$ of all A such that in this relation equal holds for $i = 1, \dots, r_B$ and $\text{rank } A \geq n - r_B$ is non-empty. In fact, if $\check{A} \in \mathcal{N}_{B,\Delta_B}$ is such that $\check{A}_{hk} = 0$ if there doesn't exist $i \in \{1, \dots, r_B\}$ such that $h, k \in \{n_i, \dots, n_{i+1} - 1\}$ and for $h', k' \in \{n_i, \dots, n_{i+1} - 1\}$ the nilpotent matrix $(\check{A}_{h'k'})$ is regular for $i = 1, \dots, r_B$ then we have $\check{A} \in \mathcal{Z}$.

If $A \in \mathcal{Z}$ we have

$$v_{n_i}^{u_{n_i}-1}, v_{n_i+1}^{u_{n_i+1}-1}, \dots, v_{n_{i+1}-1}^{u_{n_{i+1}-1}-1} \in V_{A,i}$$

for $i = 1, \dots, r_B$, which implies that $\text{Im } A$ is contained in the following subspace of K^n :

$$\langle Av_h^l : h = 1, \dots, t, l = 0, \dots, u_h - 1, (h, l) \neq (n_i, u_{n_i} - 1) \text{ for } i = 1, \dots, r_B \rangle.$$

This shows that $\mathcal{Z}$ has the required property. $\square$

3 On some automorphisms of $\mathcal{H}(n, K)$

The aim of this section is the proof of Proposition 3.5 and Proposition 3.7, from which we will get Corollary 3.8. For this purpose, we first prove some properties of the ranks of the matrices $(A^m)_{hk}$ for $A \in \mathcal{N}_B$, $m \in \mathbb{N}$ and $h, k \in \{1, \dots, t\}$.

Lemma 3.1 *Let $A \in \mathcal{N}_{B, \Delta_B}$ and let $\{i_1, \dots, i_r\} \subseteq \{1, \dots, t\}$ be maximal such that $i_1 < \dots < i_r$, $u_{i_1} - u_{i_r} \leq 1$. If $m \in \{2, \dots, r\}$ and $i, j \in \{i_1, \dots, i_r\}$ we have*

$$\text{rank}(A^m)_{ij} \leq \begin{cases} u_j - 1 & \text{if } 0 \leq j - i \leq m - 1 \\ u_j - 2 & \text{if } i - j \geq r - m + 1. \end{cases}$$

Proof. If $j - i \geq 0$ and $m \in \{2, \dots, r\}$ we have $\text{rank}(A^m)_{ij} \leq u_j - 1$ if there doesn't exist an $l \in \{1, \dots, t\}$ such that $(A^{m-1})_{il}$ and A_{lj} have the first column different from 0. If $i - j \geq 0$ and $m \in \{2, \dots, r\}$ we have $\text{rank}(A^m)_{ij} \leq u_j - 2$ if there doesn't exist an $l \in \{1, \dots, t\}$ such that either $(A^{m-1})_{il}$ has the first column different from 0 and A_{lj} has the second column different from 0 or the converse. Hence the claim is true for $m = 2$. Then by induction on m and the previous observations we get the claim. $\square$

Let s_B be the maximum of the cardinalities of the subsets $\{i_1, \dots, i_r\}$ of $\{1, \dots, t\}$ such that $i_1 < \dots < i_r$, $u_{i_1} - u_{i_r} \leq 1$.

Since if $A \in \mathcal{N}_B$ we have $(\overline{A}_{\alpha\alpha})^{s_B} = 0$ for $\alpha = 1, \dots, \hat{t}$, by Lemma 3.1 we get the following corollary.

Corollary 3.2 *If $A \in \mathcal{N}_B$, $i, j \in \{1, \dots, t\}$ and $|u_i - u_j| \leq 1$ we have $\text{rank}(A^{s_B})_{ij} \leq \min\{u_i, u_j\} - 1$. $\square$*

If X is a matrix and $m \in \mathbb{Z}$ is such that $m \leq 0$ the claims $\text{rank } X = m$ and $\text{rank } X < m$ will mean $\text{rank } X = 0$.

If $X_1 \in \mathcal{M}(p_1 \times u, K)$ and $X_2 \in \mathcal{M}(u \times p_2, K)$ have the following form:

$$X_i = \begin{pmatrix} 0 & \dots & 0 & x_1^i & x_2^i & \dots & x_{r_i}^i \\ \vdots & & & 0 & x_1^i & \ddots & \vdots \\ \vdots & & & & \ddots & \ddots & x_2^i \\ \vdots & & & & & 0 & x_1^i \\ \vdots & & & & & & 0 \\ \vdots & & & & & & \vdots \\ 0 & \dots & \dots & \dots & \dots & \dots & 0 \end{pmatrix}$$

for $i = 1, 2$, where $r_i \leq \min\{p_i, u\}$, then $X_1 X_2 \in \mathcal{M}(p_1 \times p_2, K)$ has the same form and $\text{rank}(X_1 X_2) = \text{rank} X_1 + \text{rank} X_2 - u$.

Hence, by Lemma 2.2, for $m_1, m_2 \in \mathbb{N}$ and $A \in \mathcal{N}_B$, since $A^{m_1+m_2} = A^{m_1} A^{m_2}$, we have

$$\text{rank}(A^{m_1+m_2})_{hk} \leq \max_{l \in \{1, \dots, t\}} \{\text{rank}(A^{m_1})_{hl} + \text{rank}(A^{m_2})_{lk} - u_l\}. \quad (1)$$

Let $\widehat{\mathcal{N}}_B$ be the subset of $\mathcal{N}_B$ of all A such that if $i, j \in \{1, \dots, t\}$ and $|u_i - u_j| \leq 1$ we have $\text{rank} A_{ij} \leq \min\{u_i, u_j\} - 1$.

Let $i, j \in \{1, \dots, t\}$ be such that $i \leq j$. We associate to the pair (i, j) an $h(i, j) \in \mathbb{N} \cup \{0\}$ defined as follows. If $u_i - u_j \leq 1$ we set $h(i, j) = 0$. If $u_i - u_j \geq 2$ we define $h(i, j)$ to be the unique element of $\mathbb{N}$ such that there exists $\{k_l : l = 0, \dots, h(i, j)\} \subseteq \{1, \dots, t\}$ with the following properties: $i = k_0 < k_1 < \dots < k_{h(i, j)}$, $u_{k_l} - u_{k_{l+1}} \geq 2$ and $u_{k_l} - u_{k_{l+1}-1} \leq 1$ for $l = 0, \dots, h(i, j) - 1$, $0 \leq u_{k_{h(i, j)}} - u_j \leq 1$.

Lemma 3.3 *If $A \in \widehat{\mathcal{N}}_B$, $i, j \in \{1, \dots, t\}$ and $m \in \mathbb{N}$ we have*

$$\text{rank}(A^m)_{ij} \leq \min\{u_i, u_j\} - m + h(\min\{i, j\}, \max\{i, j\}).$$

Proof. The claim is true if $m = 1$, hence we prove it by induction on m . Let $m \geq 2$.

Let $j \leq i$. By 1 we get

$$\text{rank}(A^m)_{ij} \leq \max_{l \in \{1, \dots, t\}} \{\text{rank} A_{il} + \text{rank}(A^{m-1})_{lj} - u_l\},$$

hence it is sufficient to prove that for $l = 1, \dots, t$ we have

$$\text{rank} A_{il} + \text{rank}(A^{m-1})_{lj} - u_l \leq u_i - m + h(j, i).$$

Let $l \leq j$. Then we have $\text{rank} A_{il} \leq u_i$ and equal may hold only if $u_l - u_i \geq 2$. Moreover by the inductive hypothesis we have $\text{rank}(A^{m-1})_{lj} \leq u_j - m + 1 + h(l, j)$. Hence we get

$$\text{rank} A_{il} + \text{rank}(A^{m-1})_{lj} - u_l \leq u_i + u_j - m + 1 + h(l, j) - u_l$$

where equal may hold only if $u_l - u_i \geq 2$. Since we have $u_j + h(l, j) \leq u_l$ and equal may hold only if $u_j = u_l$ we get the claim.

Let $j \leq l \leq i$. Then as above we have $\text{rank} A_{il} \leq u_i$ and equal may hold only if $u_l - u_i \geq 2$. By the inductive hypothesis we have $\text{rank}(A^{m-1})_{lj} \leq u_l - m + 1 + h(j, l)$, hence we get

$$\text{rank} A_{il} + \text{rank}(A^{m-1})_{lj} - u_l \leq u_i - m + 1 + h(j, l)$$

where equal may hold only if $u_l - u_i \geq 2$. Since $h(j, l) \leq h(j, i)$ and equal may hold only if $u_l - u_i \leq 1$, we get the claim.

Let $i \leq l$. Then we have $\text{rank } A_{il} \leq u_l$ and equal may hold only if $u_i - u_l \geq 2$. By the inductive hypothesis we have $\text{rank}(A^{m-1})_{lj} \leq u_l - m + 1 + h(j, l)$, hence we get

$$\text{rank } A_{il} + \text{rank}(A^{m-1})_{lj} - u_l \leq u_l - m + 1 + h(j, l)$$

where equal may hold only if $u_i - u_l \geq 2$. If $u_i = u_l$ then $h(j, l) = h(j, i)$ and hence we get the claim. If $u_l + 1 = u_i$ then $h(j, l) \leq h(j, i) + 1$, hence we get the claim. Let $u_i - u_l \geq 2$. We have $h(j, l) \leq h(j, i) + h(i, l) + 1$ and $u_l + h(i, l) + 1 \leq u_i$, but not in both of these equal may hold. In fact $u_l + h(i, l) + 1 = u_i$ only if $u_i - u_l = 2$ and if $u_i - u_l = 2$ we have $h(j, l) \leq h(j, i) + h(i, l)$. Hence we get the claim.

If $i \leq j$ we can repeat the same argument as above by using

$$\text{rank}(A^m)_{ij} \leq \max_{l \in \{1, \dots, t\}} \{\text{rank}(A^{m-1})_{il} + \text{rank } A_{lj} - u_l\}. \quad \square$$

Let $m \in \mathbb{N}$ and let $\mathcal{N}_m$ be the subset of $\mathcal{M}(n, K(x))$ of all Y which commutes with B and such that

$$\text{rank } Y_{ij} \leq \min\{u_i, u_j\} - m + h(\min\{i, j\}, \max\{i, j\})$$

for $i, j = 1, \dots, t$. Let q_0 and q_α , $\alpha = 1, \dots, \hat{t}$, be as in Section 2 and let $t_m = \max\{\alpha \in \{1, \dots, \hat{t}\} : u_{q_\alpha} - 2 \geq m\}$.

For $Y \in \mathcal{N}_m$, $\alpha \in \{1, \dots, t_m\}$ and $h, k \in \{q_{\alpha-1} + 1, \dots, q_\alpha\}$ let $a_{hk}^m(Y)$ be the entry of Y_{hk} of indices $(l, m + l)$, $l = 1, \dots, u_{q_\alpha} - m$, and let $d_\alpha^m(Y) = \det(a_{hk}^m(Y))$. Let $\mathcal{V}_m$ be the open subset of $\mathcal{N}_m$ of all Y such that $d_\alpha^m(Y) \neq 0$ for $\alpha = 1, \dots, t_m$. Let $\tilde{\mathcal{V}}_m$ be the subset of $\mathcal{V}_m$ of all Y such that if $i, j \in \{1, \dots, t\}$ are such that $|u_i - u_j| \geq 2$ then $Y_{ij} = 0$.

For $Y \in \mathcal{N}_m$ let $\mathcal{R}(Y)$ be the subset of K of all a such that all the entries of Y are regular in a and let $\mathcal{E}_m(Y) = \{a \in K : d_\alpha^m(Y)(a) \neq 0 \text{ for } \alpha = 1, \dots, t_m\}$.

Lemma 3.4 *If $m \in \mathbb{N}$ there exist an open subset $\mathcal{W}_m$ of $\mathcal{V}_m$ such that $\tilde{\mathcal{V}}_m \subset \mathcal{W}_m$ and a rational map $\phi_m : \mathcal{W}_m \rightarrow \mathcal{N}_m$ such that:*

- 1) *the first m columns and the last m rows of $\phi_m(Y)_{hk}$ are 0 for $Y \in \mathcal{W}_m$ and $h, k = 1, \dots, t$;*

2) if $Y \in \mathcal{W}_m$ and g is an entry of $\phi_m(Y)$ then

$$g \prod_{\alpha=1}^{t_m} d_{\alpha}^m(Y)^{2[(t-\alpha)+(t-\alpha-1)+\dots+(t-t_m)]}$$

is regular in $\mathcal{R}(Y)$;

3) $\text{rank } \phi_m(Y)(a) = \text{rank } Y(a)$ for $Y \in \mathcal{W}_m$ and $a \in \mathcal{R}(Y) \cap \mathcal{E}_m(Y)$.

Proof. If $\alpha, \beta \in \{1, \dots, t\}$ are such that $u_{q_{\alpha}} - u_{q_{\beta}} \geq 2$ and $Y \in \mathcal{V}_m$, for $i = q_{\alpha-1} + 1, \dots, q_{\alpha}$ and $r = 1, \dots, h(q_{\alpha}, q_{\beta})$ let $C^i(Y, \beta, r)$ be the submatrix of $Y^{(i)}$ having as columns the columns of $Y^{(i)}$ of indices $m+1, \dots, u_{q_{\beta}} + r$ and let $C_i(Y, \beta, r)$ be the submatrix of $Y^{(i)}$ having as rows the rows of $Y^{(i)}$ of indices $u_{q_{\alpha}} - u_{q_{\beta}} - r + 1, \dots, u_{q_{\alpha}} - m$. Let $r \in \{1, \dots, h(q_{\alpha}, q_{\beta})\}$ and let $j \in \{q_{\beta-1} + 1, \dots, q_{\beta}\}$. Let $Y \in \mathcal{V}_m$ be such that

$$\text{rank } Y_{ij}, \text{ rank } Y_{ji} \leq u_{q_{\beta}} - m + r \quad (2)$$

$$\text{rank } Y_{li}, \text{ rank } Y_{il} \leq u_{q_{\alpha}} - m \quad (3)$$

for $i = q_{\alpha-1} + 1, \dots, q_{\alpha}$ and $l = 1, \dots, q_{\alpha-1}$. Since $d_{\alpha}^m(Y) \neq 0$, it is possible to add to the submatrix of $Y^{(j)}$ whose columns are the columns of $Y^{(j)}$ of indices $m - r + 1, \dots, u_{q_{\beta}}$ a suitable linear combination over $K(x)$ of the matrices $C^i(Y, \beta, r)$, $i = q_{\alpha-1} + 1, \dots, q_{\alpha}$, getting a matrix Y' which commutes with B and such that $\text{rank}(Y')_{ij} \leq u_{q_{\beta}} - m + r - 1$ for $i = q_{\alpha-1} + 1, \dots, q_{\alpha}$. The coefficients of this linear combination are a solution of a Cramer system whose coefficients are entries of Y and whose matrix has $d_{\alpha}^m(Y)$ as determinant. Hence if g is an entry of Y' we have that $gd_{\alpha}^m(Y)$ is regular in $\mathcal{R}(Y)$.

We claim that $Y' \in \mathcal{N}_m$. In fact, it is sufficient to prove that for $i = q_{\alpha-1} + 1, \dots, q_{\alpha}$ and $l = q_{\alpha} + 1, \dots, t$ we have

$$\text{rank } C^i(Y, \beta, r)_{(l)} \leq \min\{u_l, u_{q_{\beta}}\} - m + h(\min\{l, q_{\beta}\}, \max\{l, q_{\beta}\}).$$

We have

$$\begin{aligned} \text{rank } C^i(Y, \beta, r)_{(l)} &\leq u_l - m + h(q_{\alpha}, l) - u_{q_{\alpha}} + u_{q_{\beta}} + r \leq \\ &\leq u_l - m + h(q_{\alpha}, l) - u_{q_{\alpha}} + u_{q_{\beta}} + h(q_{\alpha}, q_{\beta}). \end{aligned}$$

If $l \geq q_{\beta}$ then $h(q_{\alpha}, l) \leq h(q_{\alpha}, q_{\beta}) + h(q_{\beta}, l) + 1$ and $2h(q_{\alpha}, q_{\beta}) \leq u_{q_{\alpha}} - u_{q_{\beta}}$, but not in both of these equal may hold. In fact, if $2h(q_{\alpha}, q_{\beta}) = u_{q_{\alpha}} - u_{q_{\beta}}$ then there exists $\{k_{l'} : l' = 0, \dots, h(q_{\alpha}, q_{\beta})\}$ such that $u_{k_{l'}} - u_{k_{l'+1}} = 2$ for $l' = 0, \dots, h(q_{\alpha}, q_{\beta}) - 1$ and $k_0 = q_{\alpha}$, $k_{h(q_{\alpha}, q_{\beta})} = q_{\beta}$ and then $h(q_{\alpha}, l) =$

$h(q_\alpha, q_\beta) + h(q_\beta, l)$. If $l \leq q_\beta$ we may use the same argument by reversing the role of q_β and l . This proves the claim.

If $Y \in \tilde{\mathcal{V}}_m$ the coefficients of the above linear combination of matrices are 0. Hence there exists an open subset of $\mathcal{V}_m$ which contains $\tilde{\mathcal{V}}_m$ such that if Y belongs to it we get $Y' \in \mathcal{V}_m$. We could repeat a similar argument for the rows; in this case we would add to the submatrix of $Y_{(j)}$ whose rows are the rows of $Y_{(j)}$ of indices $1, \dots, u_{q_\beta} - m + r$ a suitable linear combination over $K(x)$ of the matrices $C_i(Y, \beta, r)$, $i = q_{\alpha-1} + 1, \dots, q_\alpha$. Let $O(\alpha, \beta, r)$ be the operation on the rows and columns of an element Y of $\mathcal{V}_m$ which satisfies 2 and 3 for $i = q_{\alpha-1} + 1, \dots, q_\alpha$, $l = 1, \dots, q_{\alpha-1}$ and $j = q_{\beta-1} + 1, \dots, q_\beta$ which consists on applying the previous operation on the columns and then the corresponding operation on the rows for $j = q_{\beta-1} + 1, \dots, q_\beta$. For $\alpha, \beta \in \{1, \dots, \hat{t}\}$ we define $O(\alpha, \beta, 0)$ to be the operation which doesn't change the given matrix.

If $\alpha, \beta \in \{1, \dots, \hat{t}\}$ and $u_{q_\alpha} - u_{q_\beta} \geq 2$ then for $l = q_\alpha + 1, \dots, t$, $i = q_{\alpha-1} + 1, \dots, q_\alpha$ and $r = 1, \dots, h(q_\alpha, q_\beta)$ we also have

$$\text{rank } C^i(Y, \beta, r)_{(l)}, \text{ rank } C_i(Y, \beta, r)^{(l)} \leq u_{q_\beta} - m + r - 1$$

since $u_l + h(q_\alpha, l) + 1 \leq u_{q_\alpha}$. Let $\mathcal{W}_m$ be the open subset of $\mathcal{V}_m$ of all Y such that it is possible to apply to Y the following operations in the order in which they are written:

$$O(\gamma, \beta, h(q_\alpha, q_\beta)), \quad \gamma = \alpha, \alpha - 1, \dots, 1, \quad \beta = \alpha + 1, \dots, \hat{t}, \quad \alpha = 1, \dots, t_m.$$

Let $\phi_m : \mathcal{W}_m \rightarrow \mathcal{N}_m$ be the map which associates to $Y \in \mathcal{W}_m$ the matrix obtained by applying to Y the previous operations in the given order. Then ϕ_m has the required properties. $\square$

The previous lemma allows us to prove the main results of this section.

Proposition 3.5 *If $A \in \mathcal{N}_B$ and $m \in \mathbb{N}$ we have $\text{rank}(A^{s_B})^m \leq \text{rank } B^m$ and $\text{rank}(B + xA^{s_B})^m \leq \text{rank } B^m$.*

Proof. The first claim is a consequence of the second one, hence let us prove the second claim. It is sufficient to prove that there exists a non-empty open subset of $\mathcal{N}_B$ such that if A belongs to it we have $\text{rank}(B + xA^{s_B})^m \leq \text{rank } B^m$. By Corollary 3.2 and Lemma 3.3 we get that $(B + xA^{s_B})^m \in \mathcal{N}_m$ for any $A \in \mathcal{N}_B$. Moreover if $A = 0$ we have $(B + xA^{s_B})^m \in \tilde{\mathcal{V}}_m$. Hence by 1) and 3) of Lemma 3.4 we get the claim. $\square$

Let F be a field and for $p \in \mathbb{N} \cup \{0\}$ let $F[x]_p = \{g \in F[x] : \deg g \leq p\}$.

Lemma 3.6 *i) If $p_1, p_2 \in \mathbb{N} \cup \{0\}$ the subset of $F[x]_{p_1} \times F[x]_{p_2}$ of all (g_1, g_2) such that $g_2 \neq 0$ and $g_1 \notin \sqrt{(g_2)}$ is open and if $p_2 \in \mathbb{N}$ it is non-empty.*

ii) If $p_1, \dots, p_r \in \mathbb{N} \cup \{0\}$, $r > 1$ and F is algebraically closed the subset of $F[x]_{p_1} \times \dots \times F[x]_{p_r}$ of all $(g_1, \dots, g_r)$ such that there exists $i \in \{1, \dots, r\}$ such that $\deg g_i = p_i$ and $1 \in (g_1, \dots, g_r)$ is open and non-empty.

Proof.

i) The claim is obvious if $p_1 = 0$ or $p_2 = 0$, hence we assume $p_1, p_2 \in \mathbb{N}$. If $(g_1, g_2) \in F[x]_{p_1} \times F[x]_{p_2}$ we have $g_1 \in \sqrt{(g_2)}$ if and only if $(g_1)^{p_2} \in (g_2)$. If $g_2 \neq 0$ this happens if and only if there exists $[a, h] \in \mathbb{P}(F \times F[x]_{p_1 p_2})$ such that $a(g_1)^{p_2} = hg_2$, which shows the claim.

ii) Let x, y be homogeneous coordinates of $\mathbb{P}^1(F)$. For $p \in \mathbb{N} \cup \{0\}$ and $g = a_0 + a_1 x + \dots + a_p x^p \in F[x]_p$ let $g_0 = a_0 y^p + a_1 x y^{p-1} + \dots + a_p x^p \in F[x, y]$. If $(g_1, \dots, g_r) \in F[x]_{p_1} \times \dots \times F[x]_{p_r}$ and there exists $i \in \{1, \dots, r\}$ such that $\deg g_i = p_i$ then $1 \in (g_1, \dots, g_r)$ if and only if there doesn't exist $[\bar{x}, \bar{y}] \in \mathbb{P}^1(F)$ such that $(g_j)_0(\bar{x}, \bar{y}) = 0$ for $j = 1, \dots, r$, which shows the claim. $\square$

Proposition 3.7 *Let $\{i_1, \dots, i_{s_B}\} \subseteq \{1, \dots, t\}$ be such that $i_1 < \dots < i_{s_B}$, $u_{i_1} - u_{i_{s_B}} \leq 1$ and $u_{i_1} \geq 2$. If $A \in \mathcal{N}_B$ is such that at least one of the determinants of the minors of $(B + xA^{s_B})^{u_{i_1}-1}$ of order $\text{rank } B^{u_{i_1}-1}$ has the maximum possible degree there exists $a(A) \in K$ such that*

$$a) \text{ rank}(B + a(A)A^{s_B})^{u_{i_1}-1} < \text{rank } B^{u_{i_1}-1}.$$

Moreover there exists a non-empty open subset $\mathcal{U}_{i_1}$ of $\mathcal{N}_B$ such that if $A \in \mathcal{U}_{i_1}$ there exists $a(A) \in K$ such that a) and the following conditions hold:

$$b) \text{ rank}(B + a(A)A^{s_B}) \geq \text{rank } B - 1;$$

$$c) \text{ rank}(B + a(A)A^{s_B})^l = \text{rank } B^l \text{ for } l = u_{i_1}, \dots, u_1.$$

Proof. By Corollary 3.2 and Lemma 3.3 we have $(B + xA^{s_B})^{u_{i_1}-1} \in \mathcal{N}_{u_{i_1}-1}$ for any $A \in \mathcal{N}_B$. Let $\mathcal{W}_{u_{i_1}-1}$ and $\phi_{u_{i_1}-1}$ be as in Lemma 3.4 and let $\overline{\mathcal{W}}_{i_1}$ be the open subset of $\mathcal{N}_B$ of all A such that $(B + xA^{s_B})^{u_{i_1}-1} \in \mathcal{W}_{u_{i_1}-1}$. For $A \in \overline{\mathcal{W}}_{i_1}$ let $d_{i_1}(A)$ be the determinant of the submatrix of $(\phi_{u_{i_1}-1}((B + xA^{s_B})^{u_{i_1}-1})_{hk})$, $h, k = 1, \dots, t$, obtained by taking the first

$u_h - (u_{i_1} - 1)$ rows and the last $u_k - (u_{i_1} - 1)$ columns of $\phi_{u_{i_1}-1}((B + xA^{s_B})^{u_{i_1}-1})_{hk}$ for $h, k = 1, \dots, \max\{i \in \{1, \dots, t\} : u_i \geq u_{i_1}\}$. Let $m_{i_1} = \sum_{i=1}^t \max\{u_i - (u_{i_1} - 1), 0\}$ and let

$$d'_{i_1}(A) = d_{i_1}(A) \prod_{\alpha=1}^{t_{u_{i_1}-1}} d_{\alpha}^{u_{i_1}-1} ((B + xA^{s_B})^{u_{i_1}-1})^{2m_{i_1}[(\hat{t}-\alpha)+(\hat{t}-\alpha-1)+\dots+(\hat{t}-t_{u_{i_1}-1})]}.$$

By 2) of Lemma 3.4 we have $d'_{i_1}(A) \in K[x]$. We claim that the subset $\overline{\mathcal{U}}_{i_1}$ of $\overline{\mathcal{W}}_{i_1}$ of all A such that $d'_{i_1}(A) \neq 0$ and

$$\prod_{\alpha=1}^{t_{u_{i_1}-1}} d_{\alpha}^{u_{i_1}-1} ((B + xA^{s_B})^{u_{i_1}-1}) \notin \sqrt{(d'_{i_1}(A))}$$

is non-empty. In fact, let $\check{A} \in \mathcal{N}_B$ be such that $\check{A}_{hk} = 0$ if $h \notin \{i_1, \dots, i_{s_B}\}$ or $k \notin \{i_1, \dots, i_{s_B}\}$ and for $h', k' = i_1, \dots, i_{s_B}$ we have $(B_{h'k'}) = \check{a}_1(\check{A}_{h'k'})^{s_B} + \check{a}_2(\check{A}_{h'k'})^{s_B+1}$ where $\check{a}_1, \check{a}_2 \in K \setminus \{0\}$. We have $d_{\alpha}^{u_{i_1}-1}((B + x\check{A}^{s_B})^{u_{i_1}-1}) = 1$ for $\alpha = 1, \dots, t_{u_{i_1}-1}$. Since $(B + x\check{A}^{s_B})^{u_{i_1}-1} \in \tilde{\mathcal{V}}_{u_{i_1}-1}$ we have $\check{A} \in \overline{\mathcal{W}}_{i_1}$. If we set $a(\check{A}) = -\check{a}_1$ then $\check{A}$ satisfies a) and $-\check{a}_1$ is the only element of K with this property. Hence by 3) of Lemma 3.4 we have $\deg d'_{i_1}(\check{A}) \geq 1$ and then $\check{A} \in \overline{\mathcal{U}}_{i_1}$.

By i) of Lemma 3.6 the subset $\overline{\mathcal{U}}_{i_1}$ is open in $\mathcal{N}_B$. If $A \in \overline{\mathcal{U}}_{i_1}$ then all the determinants of the minors of $(B + xA^{s_B})^{u_{i_1}-1}$ of order rank $B^{u_{i_1}-1}$ have a common root, hence by ii) of Lemma 3.6 we get that if $A \in \mathcal{N}_B$ is such that at least one of these determinants has the maximum possible degree then they have a common root. This proves the first claim.

If we set $a(\check{A}) = -\check{a}_1$ then $\check{A}$ satisfies also b) and c). Hence the subset $\mathcal{U}_{i_1}$ of $\overline{\mathcal{U}}_{i_1}$ of all A such that if $\text{rank } B > 1$ there exists a minor of $B + xA^{s_B}$ of order rank $B - 1$ and determinant $d(A)$ and, if $i_1 > 1$, there exists a minor of $(B + xA^{s_B})^l$ of order rank B^l and determinant $d^l(A)$ for $l = u_{i_1}, \dots, u_1 - 1$ such that if $\text{rank } B > 1$ we have

$$d(A) \prod_{\alpha=1}^{t_{u_{i_1}-1}} d_{\alpha}^{u_{i_1}-1} ((B + xA^{s_B})^{u_{i_1}-1}) \notin \sqrt{(d'_{i_1}(A))}$$

and if $i_1 > 1$ we have

$$d(A) \prod_{l=u_{i_1}}^{u_1-1} d^l(A) \prod_{\alpha=1}^{t_{u_{i_1}-1}} d_{\alpha}^{u_{i_1}-1} ((B + xA^{s_B})^{u_{i_1}-1}) \notin \sqrt{(d'_{i_1}(A))}$$

is non-empty. Hence by *i*) of Lemma 3.6 and Proposition 3.5 we get the second claim. $\square$

Corollary 3.8 *Let $\mathcal{A}$ be an open subset of $\mathcal{H}(n, K)$, let $(A, B) \in \mathcal{A}$ and let $\{i_1, \dots, i_{s_B}\} \subseteq \{1, \dots, t\}$ be such that $u_{i_1} - u_{i_{s_B}} \leq 1$. There exist $(A', B') \in \mathcal{A}$ and $f \in K[x]$ such that $x \mid f$, $B' + f(A')$ has $s_{B'+f(A')}$ Jordan blocks of order 1 and if we denote by $u'_1, \dots, u'_{t'}$ the orders of the Jordan blocks of $B' + f(A')$ we have $t' \geq t$ and $u'_i = u_i$ for $i = 1, \dots, i_1 - 1$.*

Proof. Let $\mathcal{U}_{i_1}$ be as in Proposition 3.7, let $A_1 \in \mathcal{U}_{i_1}$ be such that $(A_1, B) \in \mathcal{A}$ and let $a(A_1)$ be as in Proposition 3.7. We set $a_1 = a(A_1)$ and $s_1 = s_B$. We now set B to be the matrix $B + a_1(A_1)^{s_1}$. If it is possible, let us choose a subset $\{i_1, \dots, i_{s_B}\}$ of $\{1, \dots, t\}$ maximal such that $u_{i_1} - u_{i_{s_B}} \leq 1$ and $u_{i_1} > 1$ and let $\mathcal{U}_{i_1}$ be as in Proposition 3.7. Let $A_2 \in \mathcal{U}_{i_1}$ be such that $(A_2, B - a_1(A_1)^{s_1}) \in \mathcal{A}$ and let $a(A_2)$ be as in Proposition 3.7. We set $a_2 = a(A_2)$ and $s_2 = s_B$. We now set B to be the matrix $B + a_2(A_2)^{s_2}$. If it is possible, let us repeat the previous argument, finding a suitable matrix A_3 such that $(A_3, B - a_1(A_3)^{s_1} - a_2(A_3)^{s_2}) \in \mathcal{A}$. If we repeat this argument, by Proposition 2.4 at the end we get a matrix $B' \in \mathcal{N}(n, K)$ and a matrix $A' \in \mathcal{N}_{B'}$ which have the required properties. $\square$

Proposition 3.9 *Let $u_1 - u_2 = 2$ and $u_i - u_{i+1} \geq 2$ for $i = 2, \dots, t - 1$. There exists a non-empty open subset of $\mathcal{N}_B$ such that if A belongs to it there exists $a(A) \in K$ such that $\text{ind}(B + a(A)A) < u_1$, $\text{rank}(B + a(A)A) = n - t$.*

Proof. If $A \in \mathcal{N}_B$ by Lemma 2.3 and Lemma 3.3 the only entry of the matrix which represents $(B + xA)^{u_1-1}$ with respect to Δ_B which may be different from 0 is the entry of indices $(1, u_1)$; let us denote it by $h(A)$.

By *i*) of Lemma 3.6 it is sufficient to prove that the subset $\mathcal{K}$ of $\mathcal{N}_B$ of all A such that $h(A) \neq 0$ and there exists a minor M of $B + xA$ of order $n - t$ such that $\det M \notin \sqrt{(h(A))}$ is non-empty.

If $X \in \mathcal{M}(n, K(x, y))$ let $(X_{12})'$ be the submatrix of X_{12} obtained by taking the first u_2 rows and let $(X_{21})'$ be the submatrix of X_{21} obtained by taking the last u_2 columns.

Let $\check{A} \in \mathcal{N}(n, K(y))$ be such that $(\check{A}_{12})', (\check{A}_{21})' = I_{u_2}$, $\check{A}_{22} = yJ_{u_2}$, $\check{A}_{ij} = 0$ if $i, j \in \{1, \dots, t\}$ and $(i, j) \neq (1, 2), (2, 1), (2, 2)$; then $[\check{A}, B] = 0$.

Let us consider the following matrix of $\mathcal{M}(2, K(x, y))$:

$$H = \begin{pmatrix} x & 1 \\ 1 & x + y \end{pmatrix}$$

and for $k = 1, \dots, u_1 - 1$ let $H^k = (h_{ij}^k)$, $i, j = 1, 2$. Then we have $((\check{A} + xB)^k)_{ii} = h_{ii}^k(J_{u_i})^k$, $((\check{A} + xB)^k)_{ij}' = h_{ij}^k(J_{u_2})^{k-1}$ for $i = 1, 2$, $j \in$

$\{1, 2\} \setminus \{i\}$ and $k = 1, \dots, u_1 - 1$. For $k = 1, \dots, u_1 - 1$ and $i, j = 1, 2$ let us regard $h_{ij}^k(x, 0)$ and $h_{ij}^k(0, y)$ as polynomials over $K(x)$ and $K(y)$ respectively. By induction on k , we get that $h_{11}^k(x, 0)$ is monic of degree k , $h_{11}^k(0, y)$ is monic of degree $k - 2$, $h_{12}^k(0, y)$ and $h_{21}^k(0, y)$ are monic of degree $k - 1$, $h_{22}^k(0, y)$ is monic of degree k for $k = 2, \dots, u_1 - 1$.

If $a, b \in K$ we have $\text{rank}(\check{A}(b) + aB) < n - t$ if and only if either $t > 2$ and $a = 0$ or $\det H(a, b) = 0$.

We claim that there exists a non-empty open subset K' of K such that if b belongs to it we have

$$(\det H(x, b), h_{11}^{u_1-1}(x, b)) = 1.$$

Since $\det H = x^2 + yx - 1$ if $\text{char } K \neq 2$ this follows from the formula of the roots of $\det H$ as polynomial over $K(y)$. Let $\text{char } K = 2$. By *ii*) of Lemma 3.6 it is sufficient to prove that

$$(\det H(x, 1), h_{11}^{u_1-1}(x, 1)) = 1.$$

We have $\det H(x, 1) = x^2 + x + 1$, hence the roots of $\det H(x, 1)$ are the 3-th roots of 1 different from 1. If, for $k = 1, \dots, u_1 - 1$, we denote by $\tilde{H}_k$ the matrix over $K(x)$ obtained by substituting in $H(x, 1)^k$ the exponent of any power of x with the remainder of the division of it by 3, we have

$$\tilde{H}_2 = \begin{pmatrix} x^2 + 1 & 1 \\ 1 & x^2 \end{pmatrix}, \quad \tilde{H}_3 = \begin{pmatrix} x & x^2 + x \\ x^2 + x & x^2 \end{pmatrix}, \quad \tilde{H}_4 = \tilde{H}_1.$$

Hence $h_{11}^k(x, 1)$, for $k = 1, \dots, u_1 - 1$, hasn't a root which is a 3-th root of 1 different from 1, which proves the previous claim if $\text{char } K = 2$.

Let K'' be the non-empty open subset of K of all b such that $h_{11}^{u_1-1}(0, b) \neq 0$. Let $\bar{b} \in K' \cap K''$ and let $\bar{a} \in K \setminus \{0\}$ be such that $h_{11}^{u_1-1}(\bar{a}, \bar{b}) = 0$. Then $\text{ind}(B + \bar{a}^{-1}\check{A}(\bar{b})) < u_1$, $\text{rank}(B + \bar{a}^{-1}\check{A}(\bar{b})) = n - t$. Since $h_{11}^{u_1-1}(x, \bar{b}) \neq 0$ we have $h(\check{A}(\bar{b})) \neq 0$. Hence $\check{A}(\bar{b}) \in \mathcal{K}$. $\square$

4 Proof of the irreducibility of $\mathcal{H}(n, K)$

For $q \in \mathbb{Q}$ let $[q] = \min\{q' \in \mathbb{Z} : q' \geq q\}$. Let

$$\tilde{\mathcal{H}}(n, K) = \{(A, B) \in \mathcal{H}(n, K) : \text{rank } A \geq n - 2\}.$$

For $m \in \left\{3, \dots, \left\lceil \frac{n}{2} \right\rceil\right\}$ let

$$\mathcal{H}_m(n, K) = \{(A, B) \in \mathcal{H}(n, K) : \text{ind } A \leq n - m + 3\}$$

and let

$$\tilde{\mathcal{H}}_m(n, K) = \left\{ (A, B) \in \tilde{\mathcal{H}}(n, K) : \frac{n}{2} + 1 < \text{ind } A \leq n - m + 3 \right\}.$$

We first prove the irreducibility of the variety $\tilde{\mathcal{H}}_m(n, K)$ for $m = 3, \dots, \left\lfloor \frac{n}{2} \right\rfloor$ under some conditions on the characteristic of K .

For $l \in \left\{ 3, \dots, \left\lfloor \frac{n}{2} \right\rfloor \right\}$ let

$$J_{l,n} = \begin{pmatrix} J_l & 0 \\ 0 & 0 \end{pmatrix}.$$

Lemma 4.1 *Let $\mathcal{A}$ be an open subset of $\tilde{\mathcal{H}}(n, K)$, let $(A, B) \in \mathcal{A}$ and let $\frac{n}{2} + 1 < \text{ind } A < n$. There exist $(A', B') \in \mathcal{A}$ and $f \in K[x]$ such that $x \mid f$ and $B' + f(A')$ is conjugate to $J_{n-\text{ind } A+2,n}$.*

Proof. Since $\{A\} \times \mathcal{N}_A$ is irreducible, by Proposition 3.5 we may choose B such that A and B are conjugate. We denote by u_1 and u_2 , where $u_1 - u_2 > 2$, the orders of their Jordan blocks.

Let Δ_A be a basis of K^n such that A with respect to it is in Jordan canonical form and for $X \in \mathcal{N}_A$ let $(X_{ij}^{(A)})$, $i, j = 1, 2$ be the block matrix which represents X with respect to Δ_A , where $X_{ij}^{(A)}$ is an $u_i \times u_j$ matrix for $i = 1, 2$. By Lemma 2.2 there exists $\tilde{f} \in K[x]$ such that $x \mid \tilde{f}$ and $(B + \tilde{f}(A))_{11}^{(A)}$ is 0. Then $\text{rank}(B + \tilde{f}(A)) \leq n - 3$. If we choose B in a suitable open subset of $\mathcal{N}_A$ we have

$$\text{rank}((B + \tilde{f}(A))^h)_{12}^{(A)}, \text{rank}((B + \tilde{f}(A))^h)_{21}^{(A)} = u_2 + 1 - h,$$

$$\text{rank}((B + \tilde{f}(A))^h)_{22}^{(A)} = u_2 - h$$

and

$$\text{rank}((B + \tilde{f}(A))^h)_{11}^{(A)} = u_2 + 2 - h$$

for $h = 2, \dots, u_2 + 1$. Hence the difference between the orders of the first two Jordan blocks of $B + \tilde{f}(A)$ is greater than 1. If we apply Corollary 3.8 to $(A, B + \tilde{f}(A))$ and the open subset $\{(A'', B'' + \tilde{f}(A'')) : (A'', B'') \in \mathcal{A}\}$ by Proposition 2.4 we get that there exists $(A', B') \in \mathcal{A}$ and $f \in K[x]$ such that $x \mid f$ and, if we denote by $u'_1, \dots, u'_{t'}$ the orders of the Jordan blocks of $B' + f(A')$, we have $u'_1 = u_2 + 2$ and $u'_i = 1$ for $i = 2, \dots, t'$. This proves the claim. $\square$

For $l \in \left\{3, \dots, \left\lceil \frac{n}{2} \right\rceil\right\}$ and $c \in K \setminus \{0\}$ let

$$A_{l,n}(c) = \begin{pmatrix} J_l & E_{l,n}(c) \\ E'_{l,n} & J_{n-l} \end{pmatrix}$$

where

$$E_{l,n}(c) = \begin{pmatrix} c & 0 & \dots & 0 \\ 0 & \dots & \dots & 0 \\ \vdots & & & \vdots \\ 0 & \dots & \dots & 0 \end{pmatrix}, \quad E'_{l,n} = \begin{pmatrix} 0 & \dots & \dots & 0 \\ \vdots & & & \vdots \\ 0 & \dots & \dots & 0 \\ 0 & \dots & 0 & 1 \end{pmatrix}.$$

For $l \in \left\{2, \dots, \left\lceil \frac{n}{2} \right\rceil\right\}$ let

$$Z_l = \begin{cases} \{(n-1, 0), \dots, (0, 0)\} & \text{if } l = 2 \\ \{(n-l+1, 0), \dots, (l-2, 0), (l-3, 1), \\ (l-3, 0), (l-4, 1), (l-4, 0), \dots, (0, 1), (0, 0)\} & \text{if } l > 2, \end{cases}$$

which has cardinality n . We denote by $\{e_1, \dots, e_n\}$ the canonical basis of K^n .

Lemma 4.2 *For $l \in \left\{3, \dots, \left\lceil \frac{n}{2} \right\rceil\right\}$ the subset of $\mathcal{N}_{J_{l,n}}$ of all A such that $\text{rank } A = n-2$, $\text{ind } A = n-l+2$ is open and non-empty and if A belongs to it then:*

a) *there exist $c \in K \setminus \{0\}$, $\widehat{f} \in K[x]$ such that $\deg \widehat{f} \leq l-2$, $x \mid \widehat{f}$, $x^2 \nmid \widehat{f}$ and a basis of K^n such that the matrices which represent A and $J_{l,n}$ with respect to it are $A_{l,n}(c)$ and $\widehat{f}(J_{l,n})$ respectively;*

b) *if $v \in K^n$ is such that $(J_{l,n})^{l-1}v \neq 0$ then*

$$\{v, Av, \dots, A^{n-l+1}v, J_{l,n}v, \dots, (J_{l,n})^{l-2}v\}, \{A^i(J_{l,n})^jv : (i, j) \in Z_l\}$$

are linearly independent.

Proof. By Lemma 2.2 and Lemma 2.3 if $A \in \mathcal{N}_{J_{l,n}}$ there exists $g_A \in K[x]$ such that $x \mid g_A$ and the submatrix of A obtained by taking the first l rows and columns is $g_A(J_l)$. For $A \in \mathcal{N}_{J_{l,n}}$ let $A_0 = A - g_A(J_{l,n})$ and let A'_0 be the submatrix of A_0 obtained by taking the rows and the columns of indices $1, l, \dots, n$. Since the rows and the columns of A_0 of indices $2, \dots, l-1$ are

0 we have $\text{ind } A_0 = n - l + 2$ if and only if A'_0 , which is nilpotent, is regular. By Lemma 2.2 and Lemma 2.3 the subset of $\mathcal{N}_{J_{l,n}}$ of all A such that A'_0 is regular is non-empty. Since $A_0 J_{l,n} = 0$ we have $\text{ind } A = n - l + 2$ if and only if $\text{ind } A_0 = n - l + 2$. If A'_0 is regular we have $\text{rank } A = n - 2$ if and only if $x^2 \nmid g_A$. Hence the subset of $\mathcal{N}_{J_{l,n}}$ of all A such that $\text{rank } A = n - 2$ and $\text{ind } A = n - l + 2$ is open and non-empty. Let A belong to this subset. Then $\{e_1, \dots, e_l, (A_0)^{n-l}e_l, \dots, A_0 e_l\}$ is linearly independent and there exists $a_A \in K \setminus \{0\}$ such that $(A_0)^{n-l+1}e_l = a_A e_1$. Then $\{e_l, A e_l, \dots, A^{n-l+1}e_l, J_{l,n}e_l, \dots, (J_{l,n})^{l-2}e_l\}$ is linearly independent. Since $A J_{l,n} = g_A(J_{l,n})J_{l,n}$ and $x^2 \nmid g_A$ we get b).

Let $\gamma_1, \dots, \gamma_{l-1} \in K$, $\gamma_1 \neq 0$, be such that $J_{l,n} = \gamma_1 g_A(J_{l,n}) + \dots + \gamma_{l-1} g_A(J_{l,n})^{l-1}$ and let $\hat{f} = \gamma_1 x + \dots + \gamma_{l-2} x^{l-2}$. Let $e'_{l-i} = g_A(J_{l,n})^i e_l$ for $i = 1, \dots, l-1$. If $c' \neq 0$ is the coefficient of x in g_A we have $e'_1 = (c')^{l-1} e_1$ and hence if $c = a_A (c')^{1-l}$ we have $(A_0)^{n-l+1}e_l = c e'_1$. Then $\hat{f}$, c and the basis

$$\{e'_1, \dots, e'_{l-2}, e'_{l-1} + \gamma_1^{-1} \gamma_{l-1} e'_1, e_l + \gamma_1^{-1} \gamma_{l-1} c^{-1} (A_0)^{n-l} e_l, (A_0)^{n-l} e_l, \dots, A_0 e_l\}$$

have the property required in a). $\square$

For $l \in \mathbb{N} \cup \{0\}$ and $l' \in \mathbb{N}$ let

$$F(l, l') = \begin{cases} y_1 y + y_2 y^2 + \dots + y_{l'} y^{l'} & \text{if } l = 0 \\ x_1 x + x_2 x^2 + \dots + x_l x^l + y_1 y + y_2 y^2 + \dots + y_{l'} y^{l'} & \text{if } l \neq 0, \end{cases}$$

which we consider as a polynomial of $\mathbb{Z}[y_1, \dots, y_{l'}][x, y]$ if $l = 0$ and of $\mathbb{Z}[x_1, \dots, x_l, y_1, \dots, y_{l'}][x, y]$ if $l \neq 0$.

For $k \in \mathbb{N} \cup \{0\}$ let $F(l, l', k)$ be the polynomial obtained by substituting $x^i y^j$ with y^{i+j} in $F(l, l')$ for $i, j \in \mathbb{N}$. Let $f_0(l, l', k)$ be the term of degree 0 of $F(l, l', k)$ and for $h \in \mathbb{N}$ let $f_h(l, l', k)$ and $f'_h(l, l', k)$ be the coefficients of x^h and y^h respectively in $F(l, l', k)$.

Lemma 4.3 *Let $l \in \mathbb{N} \cup \{0\}$ and $l', h, k \in \mathbb{N}$.*

i) *If $l \neq 0$ we have*

$$\begin{aligned} f_h(l, l', k) &= x_1 f_{h-1}(l, l', k-1) + \dots + x_{\min\{l, h\}} f_{h-\min\{l, h\}}(l, l', k-1), \\ f'_{h+1}(l, l', k) &= x_1 f'_h(l, l', k-1) + \dots + x_{\min\{l, h\}} f'_{h+1-\min\{l, h\}}(l, l', k-1) + \\ &\quad + y_1 f'_h(l, l', k-1) + \dots + y_{\min\{l', h\}} f'_{h+1-\min\{l', h\}}(l, l', k-1) + \\ &\quad + y_1 f_h(l, l', k-1) + \dots + y_{\min\{l', h+1\}} f_{h+1-\min\{l', h+1\}}(l, l', k-1). \end{aligned}$$

- ii) Let $k \neq 1$. If $h \leq k - 1$ then $f_h(l, l', k), f'_h(l, l', k) = 0$. If $h > k - 1$ and $l \neq 0$ then $f_h(l, l', k)$ is a polynomial of $\mathbb{Z}[x_1, \dots, x_{\min\{l, h-k+1\}}]$ homogeneous of degree k and $f'_h(l, l', k)$ is a polynomial of $\mathbb{Z}[x_1, \dots, x_{\min\{l, h-k+1\}}, y_1, \dots, y_{\min\{l', h-k+1\}}]$ homogeneous of degree k .
- iii) If $p, m' \in \mathbb{N}$, p is prime, $p > k$ and m' is a coefficient of either one of $f_h(l, l', k), f'_h(l, l', k)$ or of the polynomial obtained by setting $y_1 = 1$ and $y_2, \dots, y_{l'} = 0$ in $f'_h(l, l', k)$ then $p \nmid m'$.

Proof. i) follows from the definitions. If $k \neq 1$, $j \in \mathbb{N} \cup \{0\}$ and $h - (k - 1) < j \leq h$ we have $f_0(l, l', k - 1), f_{h-j}(l, l', k - 1), f'_{h-j}(l, l', k - 1) = 0$, hence by i) and induction on k we get ii). If $l = 0$ then iii) may be proved by induction on l' using the relation

$$F(0, l', k) = \sum_{j=0}^k \binom{k}{j} (y_{l'})^{k-j} y_{l'}^{l'(k-j)} F(0, l' - 1, j).$$

If $l \neq 0$ then iii) may be proved by induction on l using the relation

$$\begin{aligned} F(l, l', k) &= \sum_{j=0}^k \binom{k}{j} (x_l)^{k-j} x_l^{l(k-j)} F(l - 1, l', j)(x, 0) + \\ &+ \sum_{j'=1}^k \binom{k}{j'} (x_l)^{k-j'} y_l^{l(k-j')} F(l - 1, l', j')(0, y). \quad \square \end{aligned}$$

For $l = 2, \dots, \left\lfloor \frac{n}{2} \right\rfloor$ let $\Phi(n, l) \in \mathcal{M}((n - l + 2) \times n, \mathbb{Z}[x_1, \dots, x_{n-l+1}])$ be such that the entry of $\Phi(n, l)$ of indices h, k is $f_{h-1}(n - l + 1, l', n - k)$ for $h = 1, \dots, n - l + 2$, $k = 1, \dots, n$, where l' is any element of $\mathbb{N}$.

For $l = 2, \dots, \left\lfloor \frac{n}{2} \right\rfloor$ let $\Psi(n, l) \in \mathcal{M}((l - 1) \times n, \mathbb{Z}[x_1, \dots, x_{l-2}, y_1, \dots, y_{l-1}])$ be such that the entry of $\Psi(n, l)$ of indices h', k is $f'_{h'}(l', l - 1, n - k)$ for $h' = 1, \dots, l - 1$, $k = 1, \dots, n$, where l' is any element of $\mathbb{N} \cup \{0\}$ such that $l' \geq l - 2$.

Examples. We have that $\Phi(7, 3) \in \mathcal{M}(6 \times 7, \mathbb{Z}[x_1, \dots, x_5])$ is the following matrix:

$$\begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & x_1 & 0 \\ 0 & 0 & 0 & 0 & (x_1)^2 & x_2 & 0 \\ 0 & 0 & 0 & (x_1)^3 & 2x_1x_2 & x_3 & 0 \\ 0 & 0 & (x_1)^4 & 3(x_1)^2x_2 & (x_2)^2 + 2x_1x_3 & x_4 & 0 \\ 0 & (x_1)^5 & 4(x_1)^3x_2 & 3(x_1)^2x_3 + 3x_1(x_2)^2 & 2x_1x_4 + 2x_2x_3 & x_5 & 0 \end{pmatrix}$$

and $\Psi(7, 4) \in \mathcal{M}(3 \times 7, \mathbb{Z}[x_1, x_2, y_1, y_2, y_3])$ is the following matrix:

$$\begin{pmatrix} 0 & 0 & 0 & 0 & 0 & y_1 & 0 \\ 0 & 0 & 0 & 0 & (y_1)^2 + 2x_1y_1 & y_2 & 0 \\ 0 & 0 & 0 & (y_1)^3 + 3x_1(y_1)^2 + 3(x_1)^2y_1 & 2y_1y_2 + 2x_1y_2 + 2x_2y_1 & y_3 & 0 \end{pmatrix}.$$

For $i = 0, \dots, n-1$ let $\Phi(n, l, i) \in \mathcal{M}((n-l+2) \times (n-i), \mathbb{Z}[x_1, \dots, x_{n-l+1-i}])$ be defined as follows: if $i > n-l+1$ it is the zero matrix, otherwise it is the matrix obtained by putting i rows more of zeros on the top of the submatrix of $\Phi(n, l)$ obtained by taking the first $n-l+2-i$ rows and the last $n-i$ columns. For $i = 0, \dots, n-1$ let $\Psi(n, l, i) \in \mathcal{M}((l-1) \times (n-i), \mathbb{Z}[x_1, \dots, x_{l-2-i}, y_1, \dots, y_{l-1-i}])$ be defined as follows: if $i > l-2$ it is the zero matrix, otherwise it is the matrix obtained by putting i rows more of zeros on the top of the submatrix of $\Psi(n, l)$ obtained by taking the first $l-1-i$ rows and the last $n-i$ columns. Let $N = \frac{n(n+1)}{2}$ and let us define the following matrix of $\mathcal{M}((n+1) \times N, \mathbb{Z}[x_1, \dots, x_{n-l+1}, y_1, \dots, y_{l-1}])$:

$$\Upsilon'(n, l) = \begin{pmatrix} \Phi(n, l, n-1) & \cdots & \Phi(n, l, 0) \\ \Psi(n, l, n-1) & \cdots & \Psi(n, l, 0) \end{pmatrix};$$

let $\Upsilon''(n, l)$ be the submatrix of $\Upsilon'(n, l)$ obtained by taking the first n rows and let $\Upsilon(n, l) \in \mathcal{M}(n \times N, \mathbb{Z}[x_1, \dots, x_{n-l+1}, y_1, \dots, y_{l-1}, z])$ be the matrix obtained by adding to the row of index $n-l+2$ of $\Upsilon''(n, l)$ the last row of

$\Upsilon'(n, l)$ multiplied by z .

Let

$$L_n = \{(i, j) \in \{0, \dots, n-1\} \times \{0, \dots, n-1\} : i + j < n\},$$

which has cardinality N , and let us consider in L_n the lexicographic order. We will consider any subset of L_n as an ordered subset.

Lemma 4.4 1) If $A \in N(n, K)$, $\alpha_1, \dots, \alpha_{n-1} \in K$, $B = \alpha_1 A + \dots + \alpha_{n-1} A^{n-1}$ and $v \in K^n$ is such that $A^{n-1}v \neq 0$ then for $(i, j) \in L_n$ the entries of the column of $\Upsilon(n, 2)(\alpha_1, \dots, \alpha_{n-2}, 0, 1, \alpha_{n-1})$ of indices (i, j) are the coordinates of $A^i B^j v$ with respect to the basis $\{v, Av, \dots, A^{n-1}v\}$.

2) If $l \in \left\{3, \dots, \left\lceil \frac{n}{2} \right\rceil\right\}$, $\alpha_1, \dots, \alpha_{n-l+1}, \beta_1, \dots, \beta_{l-2} \in K$, $c \in K \setminus \{0\}$,

$$B = \alpha_1 A_{l,n}(c) + \dots + \alpha_{n-l+1} (A_{l,n}(c))^{n-l+1} + \beta_1 J_{l,n} + \dots + \beta_{l-2} (J_{l,n})^{l-2}$$

and $v \in K^n$ is such that $(J_{l,n})^{l-1}v \neq 0$ then for $(i, j) \in L_n$ the entries of the column of $\Upsilon(n, l)(\alpha_1, \dots, \alpha_{n-l+1}, \beta_1, \dots, \beta_{l-2}, 0, c^{-1})$ of indices (i, j) are the coordinates of $(A_{l,n}(c))^i B^j v$ with respect to the basis

$$\{v, A_{l,n}(c)v, \dots, (A_{l,n}(c))^{n-l+1}v, J_{l,n}v, \dots, (J_{l,n})^{l-2}v\}.$$

Proof. By the expression of B as linear combination of powers of A we get 1). We have $A_{l,n}(c)J_{l,n} = (J_{l,n})^2$ and $(J_{l,n})^{l-1} = c^{-1}(A_{l,n}(c))^{n-l+1}$. Hence by the expression of B as linear combination of powers of $A_{l,n}(c)$ and $J_{l,n}$ we get 2). $\square$

Let $\Pi_n(L_n)$ be the set of all the subsets of L_n of cardinality n . If Ξ is an $n \times N$ matrix let L_n be the set of the indices of the columns of Ξ and for $I \in \Pi_n(L_n)$ let $\mu(I, \Xi)$ be the minor of Ξ having I as set of the indices of the columns.

Proposition 4.5 If $n \in \mathbb{N}$, $m \in \left\{3, \dots, \left\lceil \frac{n}{2} \right\rceil\right\}$ and either $\text{char } K = 0$ or $\text{char } K \geq \frac{n}{2}$ the variety $\tilde{\mathcal{H}}_m(n, K)$ is irreducible.

Proof. If $l \in \mathbb{N}$ and $\frac{n}{2} + 1 < l \leq n - m + 3$ the subset of $\tilde{\mathcal{H}}_m(n, K)$ of all (A, B) such that $\text{ind } A = l$ is irreducible, since if $A \in \mathcal{N}(n, K)$ the variety $\mathcal{N}_A$ is irreducible. Hence, by induction on $\left\lceil \frac{n}{2} \right\rceil - m$, it is sufficient to prove

that if $(\overline{A}, \overline{B}) \in \tilde{\mathcal{H}}_m(n, K)$, $\text{ind } \overline{A} = n - m + 2$ and $\mathcal{A}$ is an open subset of $\tilde{\mathcal{H}}_m(n, K)$ such that $(\overline{A}, \overline{B}) \in \mathcal{A}$ then

$$\mathcal{A} \cap \{(A, B) \in \tilde{\mathcal{H}}_m(n, K) : \text{ind } A = n - m + 3\} \neq \emptyset.$$

By Lemma 4.1 we may assume that there exists $\overline{f} \in K[x]$ such that $x \mid \overline{f}$ and $\overline{B} + \overline{f}(\overline{A})$ is conjugate to $J_{m,n}$. Since the map from $\tilde{\mathcal{H}}_m(n, K)$ into itself defined by $(A, B) \mapsto (A, B + \overline{f}(A))$ is an automorphism of $\tilde{\mathcal{H}}_m(n, K)$, we may assume that $\overline{B}$ is conjugate to $J_{m,n}$. Hence by a) of Lemma 4.2 we may assume that there exists $\overline{c} \in K \setminus \{0\}$ such that $\overline{A} = A_{m,n}(\overline{c})$ and there exist $\overline{\gamma}_1, \dots, \overline{\gamma}_{m-1} \in K$ such that $\overline{\gamma}_1 \neq 0$ and $\overline{B} = \overline{\gamma}_1 J_{m,n} + \overline{\gamma}_2 \overline{A} \overline{B} + \dots + \overline{\gamma}_{m-1} (\overline{A})^{m-2} \overline{B}$.

In the ring $K[x, y]/(x^{n-1}y)$ let $\xi = (x^{n-1}y) + x$, $\eta = (x^{n-1}y) + y$, $\eta' = \eta - \overline{\gamma}_2 \xi \eta - \dots - \overline{\gamma}_{m-1} \xi^{m-2} \eta$. We have

$$\xi^i \eta = \xi^i \eta' + \overline{\gamma}_2 \xi^{i+1} \eta + \dots + \overline{\gamma}_{\min\{m-1, n-1-i\}} \xi^{\min\{m-2+i, n-2\}} \eta$$

for $i = 1, \dots, n-2$, hence $\xi^{n-2} \eta = \xi^{n-2} \eta'$ and by reverse induction on i we may get an expression of $\xi^i \eta$ as linear combination over K of $\xi^i \eta', \dots, \xi^{n-2} \eta'$ for $i = 1, \dots, n-2$. Then the map from $\tilde{\mathcal{H}}_m(n, K)$ into itself defined by

$$(A, B) \mapsto (A, (\overline{\gamma}_1)^{-1} (B - \overline{\gamma}_2 AB - \dots - \overline{\gamma}_{m-1} A^{m-2} B))$$

is also an automorphism of $\tilde{\mathcal{H}}_m(n, K)$. Hence we may assume $\overline{B} = J_{m,n}$. Let

$$\widehat{\mathcal{H}}_m(n, K) = \{(A, B, v) \in \tilde{\mathcal{H}}_m(n, K) \times K^n : \dim \langle A^i B^j v : (i, j) \in Z_m \rangle = n\}$$

and let $\widehat{\mathcal{A}} = \{(A, B, v) \in \widehat{\mathcal{H}}_m(n, K) : (A, B) \in \mathcal{A}\}$. Let $\overline{v} \in K^n$ be such that $(J_{m,n})^{m-1} \overline{v} \neq 0$. By b) of Lemma 4.2 we have $(A_{m,n}(\overline{c}), J_{m,n}, \overline{v}) \in \widehat{\mathcal{A}}$. It is sufficient to prove that

$$\widehat{\mathcal{A}} \cap \{(A, B, v) \in \widehat{\mathcal{H}}_m(n, K) : \text{ind } A = n - m + 3\} \neq \emptyset.$$

The group $\text{GL}(n, K)$ acts on $\widehat{\mathcal{H}}_m(n, K)$ by the relation

$$G \cdot (A, B, v) = (G^{-1}AG, G^{-1}BG, G^{-1}v)$$

for $(A, B, v) \in \widehat{\mathcal{H}}_m(n, K)$ and $G \in \text{GL}(n, K)$. We may assume that $\widehat{\mathcal{A}}$ is stable with respect to this action.

Let $\mathcal{G}(N-n, K)$ be the grassmannian of all the subspaces of dimension $N-n$

of the N -dimensional vector space $K[x, y]/(x, y)^n$ over K . Let $\mathcal{F}_m(N-n, K)$ be the open subset of $\mathcal{G}(N-n, K)$ of all V such that $V \cap \langle (x, y)^n + x^i y^j : (i, j) \in Z_m \rangle = \{0\}$ and let $\mathcal{I}_m(N-n, K)$ be the subset of $\mathcal{F}_m(N-n, K)$ of all V which are ideals of the ring $K[x, y]/(x, y)^n$.

Let

$$\zeta_m : \widehat{\mathcal{H}}_m(n, K) \rightarrow \mathcal{G}(N-n, K)$$

be the map which associates to $(A, B, v) \in \widehat{\mathcal{H}}_m(n, K)$ the kernel of the homomorphism from $K[x, y]/(x, y)^n$ onto K^n defined by $(x, y)^n + g \mapsto g(A, B)v$ for $g \in K[x, y]$. If $(A, B, v) \in \widehat{\mathcal{H}}_m(n, K)$ all the elements of the orbit of (A, B, v) have the same image under ζ_m and the matrix having as columns the vectors $A^i B^j v$ for $(i, j) \in L_n$ is the matrix of the coefficients of a system of n linear homogeneous equations for $\zeta_m((A, B, v))$. Hence ζ_m is a morphism. Moreover $\zeta_m(\widehat{\mathcal{H}}_m(n, K)) \subseteq \mathcal{I}_m(N-n, K)$.

For $V \in \mathcal{F}_m(N-n, K)$ let α_V and β_V be the endomorphisms of the vector space $K[x, y]/V$ over K such that $\alpha_V(V + x^i y^j) = V + x^{i+1} y^j$ and $\beta_V(V + x^i y^j) = V + x^i y^{j+1}$ for $(i, j) \in Z_m$. Let A_V and B_V be the matrices of α_V and β_V respectively with respect to the basis $\{V + x^i y^j : (i, j) \in Z_m\}$, which has the vector $V + 1$ as n -th vector. Let

$$\widehat{\zeta}_m : \mathcal{F}_m(N-n, K) \rightarrow M(n, K) \times M(n, K) \times K^n$$

be defined by $\widehat{\zeta}_m(V) = (A_V, B_V, e_n)$ for $V \in \mathcal{F}_m(N-n, K)$.

Let $V \in \mathcal{F}_m(N-n, K)$ and let Θ be the matrix of the coefficients of a system of n linear homogeneous equations for V . Let $(i', j') \in L_n$ and let $\Theta^{(i', j')}$ be the column of Θ of index (i', j') . Then the n -tuple of the coordinates of $V + x^{i'} y^{j'}$ with respect to the basis $\{V + x^i y^j : (i, j) \in Z_m\}$ is the solution of the Cramer system

$$\mu(Z_m, \Theta)X = \Theta^{(i', j')}$$

where X is the column of the unknowns. Hence $\widehat{\zeta}_m$ is a morphism.

If $V \in \mathcal{I}_m(N-n, K)$ we have $\alpha_V(V + g) = V + gx$ and $\beta_V(V + g) = V + gy$ for $g \in K[x, y]$, hence $(A_V, B_V, e_n) \in \widehat{\mathcal{H}}_m(n, K)$ and $\zeta_m((A_V, B_V, e_n)) = V$. If $(A, B, v) \in \widehat{\mathcal{H}}_m(n, K)$ and $V = \zeta_m((A, B, v))$ we have that A_V, B_V are the matrices which represent A, B respectively with respect to the basis $\{A^i B^j v : (i, j) \in Z_m\}$, hence (A, B, v) and (A_V, B_V, e_n) belong to the same orbit. Then, since

$$\zeta_m(\widehat{\mathcal{A}}) = (\widehat{\zeta}_m|_{\mathcal{I}_m(N-n, K)})^{-1}(\widehat{\mathcal{A}}),$$

we have that $\zeta_m(\widehat{\mathcal{A}})$ is an open subset of $\mathcal{I}_m(N-n, K)$.

Let $\widehat{\mathcal{H}}'_m(n, K)$ be the subset of $\widehat{\mathcal{H}}_m(n, K)$ of all (A, B, v) such that either

$m = 3$ and A is regular or $m > 3$ and there exists $G \in \text{GL}(n, K)$, $c \in K \setminus \{0\}$ and $\alpha_1, \dots, \alpha_{n-2m+4} \in K$ such that $G^{-1}AG = A_{m-1,n}(c)$ and

$$G^{-1}BG = \alpha_1 A_{m-1,n}(c) + \dots + \alpha_{n-2m+4} (A_{m-1,n}(c))^{n-2m+4} + J_{m-1,n}.$$

We want to prove that

$$\zeta_m(\widehat{\mathcal{A}}) \cap \zeta_m(\widehat{\mathcal{H}}'_m(n, K)) \neq \emptyset.$$

Let $\overline{\Upsilon}(n, m)$ be the matrix over K obtained by setting $x_1, \dots, x_{n-m+1}, y_2, \dots, y_{m-1} = 0, y_1 = 1$ and $z = \bar{c}^{-1}$ in $\Upsilon(n, m)$. Let $\widehat{\Upsilon}(n, m-1)$ be the matrix over $\mathbb{Z}[x_1, \dots, x_{n-2m+4}, z]$ obtained by setting $x_{n-2m+5}, \dots, x_{n-m+2}, y_2, \dots, y_{m-2} = 0$ and $y_1 = 1$ in $\Upsilon(n, m-1)$. By using the definition of $\Upsilon(n, m)$, $\Upsilon(n, m-1)$ and Lemma 4.3 we get the following properties for the determinants of the minors of order n of $\overline{\Upsilon}(n, m)$ and $\widehat{\Upsilon}(n, m-1)$.

We have

$$\det \mu(Z_{m-1}, \widehat{\Upsilon}(n, m-1)) = 1, \quad \det \mu(Z_m, \widehat{\Upsilon}(n, m-1)) = z.$$

For $i \in \{1, \dots, n-2m+4\}$ let $Z_{m-1,i}$ be the set obtained by substituting $(n-m+2, 0)$ with $(n-m+2-i, 1)$ in Z_{m-1} ; then we have

$$\det \mu(Z_{m-1,i}, \widehat{\Upsilon}(n, m-1)) = x_i.$$

Let $I \in \Pi_n(L_n)$. If $l \in \left\{2, \dots, \left\lfloor \frac{n}{2} \right\rfloor\right\}$ we will say that I has the property $\mathcal{P}(l)$ if there exist $h_i \in \{0, \dots, i-1\}$ for $i = 1, \dots, l-2$ such that

$$I = \{(n-l, 0), \dots, (0, 0)\} \cup \{(h_i, i-h_i) : i = 1, \dots, l-2\} \cup \{(h, k)\}$$

where either $(h, k) = (n-l+1, 0)$ or $h \in \{0, \dots, l-2\}$, $k = l-1-h$.

If I hasn't the property $\mathcal{P}(m-1)$ then $\det \mu(I, \widehat{\Upsilon}(n, m-1))$ hasn't a term different from 0 and of degree 0 with respect to $x_1, \dots, x_{n-2m+4}$. If I has the property $\mathcal{P}(m-1)$ there exist $j_I \in \{0, 1\}$, $f_I \in \mathbb{Z}[x_1]$ and $F_I \in \mathbb{Z}[x_1, \dots, x_{n-2m+4}]$ such that $x_1 \mid f_I$ and

$$\det \mu(I, \widehat{\Upsilon}(n, m-1)) = z^{j_I} (1 + f_I) + F_I;$$

if $I \neq Z_m, Z_{m-1}$ and either $\text{char } K = 0$ or $\text{char } K > m-1$ by ii) and iii) of Lemma 4.3 we have $f_I \neq 0$.

We have $\det \mu(I, \overline{\Upsilon}(n, m)) \neq 0$ if and only if I has the property $\mathcal{P}(m)$. Hence $\det \mu(Z_{m-1}, \overline{\Upsilon}(n, m)) = 0$ and $\det \mu(Z_{m-1,i}, \overline{\Upsilon}(n, m)) = 0$ for $i =$

$1, \dots, n - 2m + 4$.

If I has the property $\mathcal{P}(m)$ and $(n - m + 1, 0) \in I$ then I has the property $\mathcal{P}(m - 1)$ and $j_I = 1$.

Let us assume that I has the property $\mathcal{P}(m)$ and there exists $h \in \{0, \dots, m - 2\}$ such that $(h, m - 1 - h) \in I$. Let $h_{m-2} \in \{0, \dots, m - 3\}$ be such that $(h_{m-2}, m - 2 - h_{m-2}) \in I$. Let $\widehat{M}_I$ be the minor of order 2 of $\widehat{\Upsilon}(n, m - 1)$ obtained by taking the rows of indices $n - m + 2$, $n - m + 3$ and the columns of indices $(h, m - 1 - h)$, $(h_{m-2}, m - 2 - h_{m-2})$. Then there exists $g_I \in \mathbb{Z}[x_1]$ such that $x_1 \mid g_I$ and

$$\det \mu(I, \widehat{\Upsilon}(n, m - 1)) = (1 + g_I) \det \widehat{M}_I.$$

Moreover there exists $h_I \in \mathbb{Z}[x_1]$ and $H_I \in \mathbb{Z}[x_1, \dots, x_{n-2m+4}]$ such that $x_1 \mid h_I$ and the entry of $\widehat{M}_I$ of indices $(2, 2)$ is $z(1 + h_I) + H_I$. The other entries of $\widehat{M}_I$ are polynomials of $\mathbb{Z}[x_1, \dots, x_{n-2m+4}]$. Since for $h \in \{0, \dots, m - 2\}$ we have $(n - m + 1 - h) - (m - 1 - h) + 1 = n - 2m + 3$, by ii) and iii) of Lemma 4.3 if either $\text{char } K \neq 0$ or $\text{char } K > m - 1$ the entry of $\widehat{M}_I$ of indices $(1, 1)$ is an homogeneous polynomial of $\mathbb{Z}[x_1, \dots, x_{n-2m+3}]$ of degree $m - 1 - h$ if $h \neq m - 2$ and is x_{n-2m+3} if $h = m - 2$; hence $\det \mu(I, \widehat{\Upsilon}(n, m - 1))$ has degree greater than 1.

By Lemma 4.4 we have that $\overline{\Upsilon}(n, m)$ is the matrix of the coefficients of a system of n linear homogeneous equations for $\zeta_m((A_{m,n}(\overline{c}), J_{m,n}, \overline{v}))$ and if $c \in K \setminus \{0\}$, $\alpha_1, \dots, \alpha_{n-2m+4} \in K$ and $v \in K^n$ is such that $(J_{m-1,n})^{m-2}v \neq 0$ then $\widehat{\Upsilon}(n, m - 1)(\alpha_1, \dots, \alpha_{n-2m+4}, c^{-1})$ is the matrix of the coefficients a system of n linear homogeneous equations for $\zeta_m((A_{m-1,n}(c), \alpha_1 A_{m-1,n}(c) + \dots + \alpha_{n-2m+4} A_{m-1,n}(c))^{n-2m+4} + J_{m-1,n}, v))$.

Then if either $\text{char } K = 0$ or $\text{char } K > m - 1$ we get that the coordinates of $\zeta_m((A_{n,m}(\overline{c}), J_{m,n}, \overline{v}))$ as element of $\mathcal{G}(N - n, K)$ satisfy the equations of the closure in $\mathcal{G}(N - n, K)$ of $\zeta_m(\widehat{\mathcal{H}}'_m(n, K))$, and hence $\widehat{\mathcal{A}} \cap \widehat{\mathcal{H}}'_m(n, K) \neq \emptyset$. This proves the result. $\square$

Remark. There exist $n, m, p \in \mathbb{N}$ such that $m \in \left\{3, \dots, \left\lceil \frac{n}{2} \right\rceil\right\}$, p is prime, $p \leq m - 1$ and if $\text{char } K = p$ the variety $\widetilde{\mathcal{H}}_m(n, K)$ is reducible (and hence the variety $\mathcal{H}_m(n, K)$ is reducible). In fact, if $\text{char } K = 3$ and

$$I = \{(3, 0), (2, 0), (1, 0), (0, 3), (0, 2), (0, 1), (0, 0)\}$$

we have $\det \mu(I, \Upsilon(7, 4)) \neq 0$ but $\det \mu(I, \Upsilon(7, 3)) = 0$; hence by the proof of Proposition 4.5, Lemma 4.1 and Lemma 4.4 if $\text{char } K = 3$ we have that $\widetilde{\mathcal{H}}_4(7, K)$ is reducible.

Lemma 4.6 *Let $\nu : \mathcal{Y}_1 \rightarrow \mathcal{Y}_2$ be a closed surjective morphism of quasi projective varieties and let $\mathcal{Y}_2$ be irreducible. If there exists a dense subset $\tilde{\mathcal{Y}}_2$ of $\mathcal{Y}_2$ such that $\nu^{-1}(\tilde{\mathcal{Y}}_2) = \mathcal{Y}_1$ and the fibers of the elements of $\tilde{\mathcal{Y}}_2$ are all irreducible of the same dimension then $\mathcal{Y}_1$ is irreducible.*

Proof. It is a generalization of Theorem 8 of chap. I, §6 of [13]. $\square$

Now we can prove the result which is the aim of this paper.

Theorem 4.7 *If $n \in \mathbb{N}$ and either $\text{char } K = 0$ or $\text{char } K \geq \frac{n}{2}$ the variety $\mathcal{H}(n, K)$ is irreducible of dimension $n^2 - 1$.*

Proof. We prove the theorem by showing that if $(\bar{A}, \bar{B}) \in \mathcal{H}(n, K)$ and $\mathcal{A}$ is an open subset of $\mathcal{H}(n, K)$ such that $(\bar{A}, \bar{B}) \in \mathcal{A}$ there exists $(\hat{A}, \hat{B}) \in \mathcal{A}$ such that $\hat{B}$ is regular. Let $\bar{u}_1, \dots, \bar{u}_{\bar{t}}$ be the orders of the Jordan blocks of $\bar{B}$. By induction on $\bar{t}$ it is sufficient to show that there exists $(\hat{A}, \hat{B}) \in \mathcal{A}$ such that either $\hat{B}$ is regular or $\text{rank } \hat{B} > n - \bar{t}$.

If there exists $(\hat{A}, \hat{B}) \in \mathcal{A}$ such that either $\hat{A}$ is regular or $\text{rank } \hat{A} > n - \bar{t}$ then by the irreducibility of $\mathcal{N}_{\hat{A}}$ there exists $\hat{B} \in \mathcal{N}_{\hat{A}}$ such that $(\hat{A}, \hat{B}) \in \mathcal{A}$ and either $\hat{B}$ is regular or $\text{rank } \hat{B} > n - \bar{t}$.

Let $n = d_{\bar{t}}\bar{t} + r_{\bar{t}}$ where $d_{\bar{t}}, r_{\bar{t}} \in \mathbb{N} \cup \{0\}$ and $r_{\bar{t}} < \bar{t}$. Then $\text{ind } \bar{B} \geq d_{\bar{t}} + \min\{1, r_{\bar{t}}\}$. If $\text{ind } \bar{B} = d_{\bar{t}} + \min\{1, r_{\bar{t}}\}$ we have $\bar{u}_1 - \bar{u}_{\bar{t}} \leq 1$, hence by Lemma 2.4 and the irreducibility of $\mathcal{N}_{\bar{B}}$ there exists $\hat{A} \in \mathcal{N}_{\bar{B}}$ such that $\hat{A}$ is regular and $(\hat{A}, \bar{B}) \in \mathcal{A}$. Then, for a given $\bar{t}$, we may prove the claim by induction on $\text{ind } \bar{B}$.

If there exists $i \in \{1, \dots, \bar{t} - 1\}$ such that $\bar{u}_i - \bar{u}_{i+1} \leq 1$ by Proposition 2.4 and the irreducibility of $\mathcal{N}_{\bar{B}}$ there exists $\hat{A} \in \mathcal{N}_{\bar{B}}$ such that $(\hat{A}, \bar{B}) \in \mathcal{A}$ and $\text{rank } \hat{A} > n - \bar{t}$; hence we may assume $\bar{u}_i - \bar{u}_{i+1} \geq 2$ for $i = 1, \dots, \bar{t} - 1$.

Let $\bar{u}_1 - \bar{u}_2 = 2$. By Proposition 3.9 and the irreducibility of $\mathcal{N}_{\bar{B}}$ there exist $\tilde{A} \in \mathcal{N}_{\bar{B}}$ and $a(\tilde{A}) \in K$ such that $(\tilde{A}, \bar{B}) \in \mathcal{A}$, $\text{rank}(\bar{B} + a(\tilde{A})\tilde{A}) = n - \bar{t}$, $\text{ind}(\bar{B} + a(\tilde{A})\tilde{A}) < u_1$. By the inductive hypothesis applied to $(\tilde{A}, \bar{B} + a(\tilde{A})\tilde{A})$ and the open subset $\{(A'', B'' + a(\tilde{A})A'') : (A'', B'') \in \mathcal{A}\}$ there exists $(\hat{A}, \tilde{B}) \in \mathcal{A}$ such that either $\tilde{B} + a(\tilde{A})\hat{A}$ is regular or $\text{rank}(\tilde{B} + a(\tilde{A})\hat{A}) > n - \bar{t}$; hence by the irreducibility of $\mathcal{N}_{\tilde{B} + a(\tilde{A})\hat{A}}$ it is possible to choose $\hat{A}$ such that either $\hat{A}$ is regular or $\text{rank } \hat{A} > n - \bar{t}$.

Let $\bar{u}_1 - \bar{u}_2 > 2$. If $\bar{t} = 2$ the claim follows by Proposition 4.5, hence we may assume $\bar{t} > 2$. Let $\bar{n} = \bar{u}_1 + \bar{u}_2$ and let $Q \in \mathcal{N}(n - \bar{n}, K)$ be the matrix in Jordan canonical form which has $\bar{t} - 2$ Jordan blocks of orders $\bar{u}_3, \dots, \bar{u}_t$. Let $\mathcal{H}_{\bar{u}_1, \bar{u}_3}(\bar{n}, K)$ be the subvariety of $\mathcal{H}(\bar{n}, K)$ of all (C, D) such

that $\text{rank } D \geq \bar{n} - 2$, $\bar{u}_1 \leq \text{ind } D \leq \bar{n} - (\bar{u}_3 + 1)$. For $A, B \in \mathcal{M}(n, K)$ let $A = (A_{ij})$, $B = (B_{ij})$ where $i, j = 1, 2$ and $A_{11}, B_{11} \in \mathcal{M}(\bar{n}, K)$. Let $\mathcal{H}_{\bar{u}_1, \bar{u}_3, Q}(n, K)$ be the subvariety of $\mathcal{H}(n, K)$ of all (A, B) such that $B_{12}, B_{21} = 0$, $B_{22} = Q$ and $(A_{11}, B_{11}) \in \mathcal{H}_{\bar{u}_1, \bar{u}_3}(\bar{n}, K)$. Let

$$\pi_{\bar{u}_1, \bar{u}_3, Q} : \mathcal{H}_{\bar{u}_1, \bar{u}_3, Q}(n, K) \rightarrow \mathcal{H}_{\bar{u}_1, \bar{u}_3}(\bar{n}, K)$$

be defined by $\pi_{\bar{u}_1, \bar{u}_3, Q}(A, B) = (A_{11}, B_{11})$. Since $\mathcal{H}_{\bar{u}_1, \bar{u}_3, Q}(n, K)$ is homogeneous with respect to the entries of A_{12} , A_{21} , A_{22} the morphism $\pi_{\bar{u}_1, \bar{u}_3, Q}$ is closed. By Proposition 4.5 the variety $\mathcal{H}_{\bar{u}_1, \bar{u}_3}(\bar{n}, K)$ is irreducible and by Lemma 2.2 and Lemma 2.3 all the fibers of $\pi_{\bar{u}_1, \bar{u}_3, Q}$ have the same dimension. Hence by Lemma 4.6 the variety $\mathcal{H}_{\bar{u}_1, \bar{u}_3, Q}(n, K)$ is irreducible. Then there exists $(A', B') \in \mathcal{A}$ such that the Jordan blocks of B' have orders $\bar{n} - (\bar{u}_3 + 1), \bar{u}_3 + 1, \bar{u}_3, \dots, \bar{u}_{\bar{t}}$. If B' has this property there exists $\hat{A} \in \mathcal{N}_{B'}$ such that $\text{rank } \hat{A} > n - \bar{t}$ and hence we get the claim. $\square$.

By Theorem 4.7 we get a proof of the following result and an extension of it to algebraically closed fields K such that $\text{char } K \geq \frac{n}{2}$.

Corollary 4.8 (J. Briançon, 1977). *If $n \in \mathbb{N}$, $\mathcal{X}$ is an algebraic surface over an algebraically closed field K such that either $\text{char } K = 0$ or $\text{char } K \geq \frac{n}{2}$ and $P \in \mathcal{X}$ is nonsingular then $\text{Hilb}^n(\mathcal{O}_P)$ is irreducible of dimension $n - 1$.*

Proof. By [10] the quotient of $\hat{\mathcal{H}}(n, K)$ with respect to the action of $\text{GL}(n, K)$ is a variety isomorphic to $\text{Hilb}^n(\mathcal{O}_P)$. Since $\hat{\mathcal{H}}(n, K)$ is an open subset of $\mathcal{H}(n, K) \times K^n$ by Theorem 4.7 it is irreducible of dimension $n^2 - 1 + n$. Since the stabilizer of any element of $\hat{\mathcal{H}}(n, K)$ with respect to the action of $\text{GL}(n, K)$ is trivial, we get the claim. $\square$

References

- [1] V. Baranovsky, The Variety of Pairs of Commuting Nilpotent Matrices is Irreducible, *Transform. Groups* **6** n. 1 (2001) 3-8.
- [2] R. Basili, On the Irreducibility of Varieties of Commuting Matrices, *J. Pure Appl. Algebra* **149** n. 2 (2000) 107-120.
- [3] J.P. Brennan, On the Normality of Commuting Varieties of Symmetric Matrices, *Comm. Algebra* **22** (15) (1994) 6409-6415.
- [4] J. Briançon, Description de $\text{Hilb}^n \mathbb{C}\{x, y\}$, *Invent. Math.* **41** n. 1 (1977) 45-89.

- [5] F.R. Gantmacher, The Theory of Matrices, Chelsea Publishing Company, New York (1959).
- [6] M. Gerstenhaber, On Dominance and Varieties of Commuting Matrices, *Ann. Math.* **73** (1961) 324-348.
- [7] M. Granger, Géométrie des schémas de Hilbert ponctuels, *Mém. Soc. Math. Fr.* n. 8 (1983).
- [8] A.A. Iarrobino, Punctual Hilbert Schemes, *Mem. Amer. Math. Soc.* **10** n. 188 (1977).
- [9] T.S. Motzkin and O. Taussky, Pairs of Matrices with Property L II, *Trans. Amer. Math. Soc.* **80** (1955) 387-401.
- [10] H. Nakajima, Lectures on Hilbert Schemes of Points on Surfaces, University Lecture Series 18, Providence, RI: American Mathematical Society (1999).
- [11] D.I. Panyushev, The Jacobian Modules of a Representation of a Lie Algebra and Geometry of Commuting Varieties, *Compositio Math.* **94** (1994) 181-199.
- [12] R.W. Richardson, Commuting Varieties of Semisimple Lie Algebras and Algebraic Groups, *Compositio Math.* **38** (1979) 311-327.
- [13] I.R. Shafarevic, Basic Algebraic Geometry, Springer-Verlag, Berlin Heidelberg New York (1977).
- [14] H.W. Turnbull, A.C. Aitken, An Introduction to the Theory of Canonical Matrices, Dover Publications Inc., New York (1961).